

BLOW UP IN THE SOLUTIONS OF AN EQUATION DESCRIBING A TRANSVERSE MOTION OF A NONHOMOGENEOUS STRING

SALIM A. MESSAOUDI

ABSTRACT. We prove a blow up result for the equation $w_{tt} = a(w_t)\varphi(w_x)w_{xx}$, which can be taken as a model for a transverse motion of a string with a density depending on the velocity.

1. INTRODUCTION

The aim of this paper is to study the existence and nonexistence of classical solutions to the one-dimensional nonlinear equation of the form

$$(1.1) \quad w_{tt}(x, t) = a(w_t(x, t))b(w_x(x, t))w_{xx}(x, t),$$

where $x \in I$ (bounded or unbounded interval), $t \geq 0$. This equation can be regarded as a model for a transverse motion of a nonhomogeneous vibrating string, where the density is depending on the velocity w_t . By assuming that

$$(1.2) \quad a(\xi) > 0, \quad b(\xi) > 0, \quad \forall \xi \in \mathbb{R},$$

the equation (1.1) is strictly hyperbolic.

Generally, classical solutions of problems associated with (1.1) develop singularities in finite time, if the elastic response functions a and b satisfy some 'genuine' nonlinearity conditions. For $a \equiv 1$, Lax [5] and MacCamy and Mizel [9] studied the problem and showed that classical solutions break down in finite time even for smooth and small initial data. In his work, Lax assumed

1991 Mathematics Subject Classification. 35L45.

that b' does not change sign, whereas MacCamy and Mizel allowed b' to change sign. They also showed, under appropriate conditions on b , that intervals of x can exist, in which the solution must exist for all time t even though it breaks down for values x outside these intervals. For a depending on x , Messaoudi [11] discussed the Cauchy problem and established a blow up result for smooth initial data.

In the dissipative case, the situation is different. For initial data small and smooth enough, the effect of the damping term dominates the nonlinear elastic response and global solutions can be obtained (see [12]). However, for large initial data the nonlinearity in the elastic response takes over and classical solutions may develop singularities in finite time. These results have been established by several authors (see [4], [10], [14]).

It is interesting to mention that nonlinear hyperbolic systems, of which equation (1.1) with $a \equiv 1$ is a special case, have attracted the attention of many authors and several results concerning global existence and blow up have been established (see [6], [7], [8], [13]).

This work will be divided into two parts. In the first part we state, without proof, a local existence result. In the second part, we state and prove our main blow up result.

2. LOCAL EXISTENCE

In this section we state a local existence theorem. The proof is omitted since it can be easily established by either using a classical energy argument [1], or applying the nonlinear semigroup theory presented in [2]. We set

$$u(x, t) := w_t(x, t), \quad v(x, t) := w_x(x, t)$$

and substitute in (1.1) to obtain the system

$$\begin{aligned} u_t(x, t) &= a(u(x, t)) b(v(x, t)) v_x(x, t) \\ (2.1) \quad v_t(x, t) &= u_x(x, t), \quad x \in \mathbb{R}, \quad t \geq 0. \end{aligned}$$

We consider (2.1) together with the initial data

$$(2.2) \quad u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), \quad x \in \mathbb{R}.$$

In order to state the local existence result, we make the following hypotheses

(H1) a and $b \in C^2(\mathbb{R})$

(H2) $a(\xi) \geq \alpha$, $b(\xi) \geq \alpha$, $\xi \in \mathbb{R}$, $\alpha > 0$.

Proposition. *Assume that (H1), (H2) hold and let u_0, v_0 in $H^2(\mathbb{R})$ be given. Then the initial value problem (2.1), (2.2) has a unique local solution (u, v) defined on a maximal time interval $[0, T)$, T depends on (u_0, v_0) , such that*

$$(2.3) \quad u, v \in C((0, T); H^2(\mathbb{R})) \cap C^1((0, T); H^1(\mathbb{R})).$$

Remark 2.1. The Sobolev embedding theorem implies that u, v are C^1 functions on $\mathbb{R} \times [0, T)$. Hence (u, v) is a classical solution.

Remark 2.2. If a and b are C^{k+1} functions and $u_0, v_0 \in H^k(\mathbb{R})$, then $u(\cdot, t), v(\cdot, t) \in H^k(\mathbb{R})$, $k \geq 1$.

Remark 2.3. A similar result can be obtained, if (H1) and (H2) hold only in a neighbourhood of zero. In this case, we have to be careful with the choice of the initial data.

3. FORMATION OF SINGULARITIES

In this section, we state and prove our main result. We first start with establishing uniform bounds on the solution (u, v) in terms of the initial data.

Lemma. *Assume (H1), (H2) hold. Then for any $\varepsilon > 0$, there exists $\delta > 0$ such that given any u_0, v_0 in $H^2(\mathbb{R})$ satisfying*

$$(3.1) \quad |u_0(x)| < \delta, \quad |v_0(x)| < \delta, \quad \forall x \in \mathbb{R},$$

the solution (2.3) obeys

$$(3.2) \quad |u(x, t)| < \varepsilon, \quad |v(x, t)| < \varepsilon, \quad \forall x \in \mathbb{R}, \quad t \in [0, T).$$

Proof. We define the quantities

$$(3.3) \quad \begin{aligned} r(x, t) &:= A(u(x, t)) + B(v(x, t)) \\ s(x, t) &:= A(u(x, t)) - B(v(x, t)) \end{aligned}$$

where

$$(3.4) \quad A(z) = \int_0^z \sqrt{\frac{1}{a(\xi)}} d\xi, \quad B(z) = \int_0^z \sqrt{b(\xi)} d\xi,$$

and the differential operators

$$(3.5) \quad \begin{aligned} \partial_t^- &:= \frac{\partial}{\partial t} - \rho(x, t) \frac{\partial}{\partial x} \\ \partial_t^+ &:= \frac{\partial}{\partial t} + \rho(x, t) \frac{\partial}{\partial x}, \end{aligned}$$

where

$$(3.6) \quad \rho(x, t) = \sqrt{a(u(x, t))b(v(x, t))}.$$

Straightforward computations yield

$$(3.7) \quad \partial_t^- r = \partial_t^+ s = 0.$$

Therefore as long as a smooth solution continues to exist, r and s remain constant along backward and forward characteristics respectively; hence

$$(3.8) \quad \|r(\cdot, t)\|_\infty = \|r_0\|_\infty, \quad \|s(\cdot, t)\|_\infty = \|s_0\|_\infty, \quad \forall t \in [0, T],$$

where

$$\|\theta\|_\infty = \text{Sup}\{|\theta(x)|, x \in \mathbb{R}\}.$$

By using (3.3), (H1), and (H2), the lemma is established.

Now we proceed to prove the blow up result. To do this we assume that the solution continues to exist for ever ($T = \infty$) and then show that this leads to a contradiction by the fact that, at least one of the derivatives of the solution becomes infinite in the L^∞ norm in finite time.

Theorem. *Assume, in addition to (H1) and (H2), that a and b satisfy*

$$(3.9) \quad \frac{a'(0)}{\sqrt{a(0)}} + \frac{b'(0)}{b(0)\sqrt{b(0)}} > 0.$$

Then there exist initial data u_0, v_0 satisfying (2.3), for which the solution of the problem (2.1), (2.2) blows up in finite time.

Proof. We take an x -partial derivative of (3.7) to get

$$(3.10) \quad (\partial_t^- r)_x = r_{xt} - \rho r_{xx} - r_x \rho_x = 0$$

which implies

$$(3.11) \quad \partial_t^-(r_x) = r_x \rho_x = \frac{[a(u)b(v)]_x}{2\sqrt{a(u)b(v)}} r_x = \frac{a'(u)b(v)u_x + a(u)b'(v)v_x}{2\sqrt{a(u)b(v)}} r_x.$$

By using

$$(3.12) \quad r_x = \frac{u_x}{\sqrt{a(u)}} + \sqrt{b(v)} v_x, \quad s_x = \frac{u_x}{\sqrt{a(u)}} - \sqrt{b(v)} v_x$$

and substituting in (3.11) we obtain, by direct calculations,

$$(3.13) \quad \partial_t^- r_x = \frac{1}{4}\sqrt{ab} \left(\frac{a'}{\sqrt{a}} + \frac{b'}{b\sqrt{b}} \right) r_x^2 + \frac{1}{4}\sqrt{ab} \left(\frac{a'}{\sqrt{a}} - \frac{b'}{b\sqrt{b}} \right) r_x s_x.$$

To handle the last term in (3.13), we set

$$(3.14) \quad W := \alpha(u) \beta(v) r_x,$$

for α and β to be chosen suitably; thus we have

$$(3.15) \quad \begin{aligned} \partial_t^- W &= \frac{1}{4}\alpha(u) \beta(v) \sqrt{ab} \left(\frac{a'}{\sqrt{a}} + \frac{b'}{b\sqrt{b}} \right) r_x^2 \\ &+ \frac{1}{4}\alpha(u) \beta(v) \sqrt{ab} \left(\frac{a'}{\sqrt{a}} - \frac{b'}{b\sqrt{b}} \right) r_x s_x \\ &+ r_x (\alpha' \beta u_t + \alpha \beta' v_t - \sqrt{ab} \alpha' \beta u_x - \sqrt{ab} \alpha \beta' v_x). \end{aligned}$$

At this point, we choose α and β so that

$$(3.16) \quad \begin{aligned} &\frac{1}{4}\alpha\beta\sqrt{ab} \left(\frac{a'}{\sqrt{a}} - \frac{b'}{b\sqrt{b}} \right) r_x s_x + r_x (\alpha' \beta u_t + \alpha \beta' v_t) \\ &- r_x (\sqrt{ab} \alpha' \beta u_x - \sqrt{ab} \alpha \beta' v_x) = 0. \end{aligned}$$

By using (3.12) to substitute for s_x and (2.1), (2.2) to substitute for u_t and v_t we arrive, by simple computations, at

$$(3.17) \quad \begin{aligned} &\frac{1}{4}\alpha\beta\sqrt{ab} \left(\frac{a'}{\sqrt{a}} - \frac{b'}{b\sqrt{b}} \right) \left(\frac{u_x}{\sqrt{a}} - \sqrt{b} v_x \right) r_x + ((\alpha' \beta ab - \alpha \beta' \sqrt{ab}) v_x \\ &+ (\alpha \beta' - \alpha' \beta \sqrt{ab}) u_x) r_x = 0, \end{aligned}$$

which yields

$$(3.18) \quad \begin{aligned} & \frac{1}{4} \alpha \beta \sqrt{ab} \left(\frac{a'}{\sqrt{a}} - \frac{b'}{b\sqrt{b}} \right) \left(\frac{u_x}{\sqrt{a}} - \sqrt{b} v_x \right) \\ & + \sqrt{a} \left((\alpha \beta' - \alpha' \beta \sqrt{ab}) \right) \left(\frac{u_x}{\sqrt{a}} - \sqrt{b} v_x \right) = 0 \end{aligned}$$

hence

$$(3.19) \quad \frac{1}{4} \alpha \beta \sqrt{ab} \left(\frac{a'}{\sqrt{a}} - \frac{b'}{b\sqrt{b}} \right) + \sqrt{a} \left((\alpha \beta' - \alpha' \beta \sqrt{ab}) \right) = 0.$$

By using the fact that a and α depend on u only and that b and β depend on v only, we get

$$(3.20) \quad \frac{1}{4} \frac{a'}{a} = \frac{\alpha'}{\alpha}, \quad \frac{1}{4} \frac{b'}{b} = \frac{\beta'}{\beta}.$$

Direct integration then yields

$$\alpha(u) = a^{1/4}(u), \quad \beta(v) = b^{1/4}(v)$$

Consequently (3.13) is reduced to

$$(3.21) \quad \partial_t^- W = \frac{1}{4} a^{1/4} b^{1/4} \left(\frac{a'}{\sqrt{a}} + \frac{b'}{b\sqrt{b}} \right) W^2.$$

Up to this point we choose initial data small enough in L^∞ norm with derivatives satisfying

$$\frac{u'_0}{\sqrt{a(u_0)}} + \sqrt{b(v_0)} v'_0 > 0$$

so that the coefficient of the quadratic term in (3.21) remains bounded away from zero; i.e

$$\frac{1}{4} a^{1/4} b^{1/4} \left(\frac{a'}{\sqrt{a}} + \frac{b'}{b\sqrt{b}} \right) \geq \lambda > 0;$$

hence (3.21) yields

$$(3.22) \quad \partial_t^- W \geq \lambda W^2.$$

Therefore (3.22) shows that W (hence r_x) blows up in finite time $T_W < \infty$.

Remark 3.1. The blow up of r_x implies that either u_x or v_x (hence v_t or u_t) blows up in finite time, however the solution (u, v) remains bounded in the

L^∞ norm. Therefore it is possible for the solution to be continued , beyond the blow up, by shocks or any other type of weak solutions.

Remark 3.2. Similar result can be established for $a'/\sqrt{a} + b'/b\sqrt{b} < 0$. In this case we consider the evolution of s_x on the forward characteristics.

Acknowledgements : The author would like to thank KFUPM for its support.

REFERENCES

1. Dafermos, C. M. and W. J. Hrusa, Energy methods for quasilinear hyperbolic initial boundary value problems. Applications to Elastodynamics, *Arch. Rational Mech. Anal.* **87** (1985), 267–292.
2. Hughes, T. J. R. and Kato, T., and Maarsden, J. E., Well-posed quasilinear second order hyperbolic systems with applications to nonlinear elastodynamic and general relativity, *Arch. Rational Mech. Anal.* **63** (1977), 273–294.
3. Keyfitz, B. L. and H. C. Kranzer, A system of nonstrictly hyperbolic conservation laws arising in elasticity theory, *Arch. Rational Mech. Anal.* **72** (1980), 219–241.
4. Kosinsky, W., Gradient catastrophe in the solutions of nonconservative hyperbolic systems, *J. Math. Anal.* **61** (1977), 672–688.
5. Lax, P. D., Development of singularities in solutions of nonlinear hyperbolic partial differential equations, *J. Math. Physics* **5** (1964), 611–613.
6. Li Ta-Tsien, Zhou Yi, and Kong De-Xing, Global classical solutions for general quasilinear hyperbolic systems with decay initial data, *Nonlinear Analysis*, **28 # 1** (1997), 1299–1332.
7. Liu Fagui and Yiang Zejiang, Formation of singularities of singularities of solutions for Cauchy problem of quasilinear hyperbolic systems with dissipative terms, *J. Partial Diff. Eqns.* **5 # 1** (1992), 79–89.

8. Liu Tai Ping, Development of singularities in nonlinear waves for quasilinear hyperbolic partial differential equations, *J. Diff. Eqns.* **33** (19979), 92-111.
9. MacCamy, R.C. and Mizel, V. J., Existence and nonexistence in the large solutions of quasilinear wave equations, *Arch. Rational Mech. Anal.* **25** (1967), 299–320.
10. Messaoudi, S. A., Formation of singularities in heat propagation guided by second sound, *J. Diff. Eqns.*, **130** (1996), 92–99.
11. Messaoudi, S. A., Formation of singularities in solutions of a wave equation, *Applied Math. Letters* **12** (1999), 23 - 28.
12. Nishida, T., Global smooth solutions for the second order quasilinear wave equations with first order dissipation . *Unpublished note*, 1975.
13. Nishida, T., Nonlinear hyperbolic systems and related topics in fluid dynamics, *Publications Mathematiques D'orsay* **78–02**, Department de Mathematicques, Paris Sud 1978.
14. Slemrod, M., Instability of steady shearing flows in nonlinear viscoelastic fluid, *Arch. Rational Mech. Anal.* **68** (1978), 211–225.

MATHEMATICAL SCIENCES DEPARTMENT, KFUPM, DHAHRAN 31261, SAUDI ARABIA.

email: messaoud@kfupm.edu.sa

Date received December 21, 1998.