

Corrigendum to "On some spaces of lacunary convergent sequences derived by Norlund-type mean and weighted lacunary statistical convergence" [Arab J. Math. Sci. 20 (2) (2014) 250–263]

METIN BASARIR^{a,*}, SUKRAN KONCA^b

^a Department of Mathematics, Sakarya University, 54187, Sakarya, Turkey ^b Department of Mathematics, Bitlis Eren University, 13000, Bitlis, Turkey

Received 18 December 2014; received in revised form 23 December 2015; accepted 23 December 2015 Available online 28 January 2016

In the above referenced paper, we obtained a new lacunary sequence $\theta' = (P_{k_r})$ using a sequence of positive real numbers (p_k) . But we did not notice a lack of provision in the article. Throughout the paper, $H_r \to \infty$ as $r \to \infty$ should be assumed. The reason for adding such a provision is that the assumption " $P_n \to \infty$ as $n \to \infty$ " may not be enough to obtain the condition " $H_r \to \infty$ as $r \to \infty$ ".

Also, in the hypothesis of Theorem 6, the condition $I'_r \subset I_r$ should be added, whereas the condition $I_r \subset I'_r$ should be added in the hypothesis of Theorem 7 and Corollary 3.

DOI of original article: http://dx.doi.org/10.1016/j.ajmsc.2013.09.002. * Corresponding author.

E-mail address: basarir@sakarya.edu.tr (M. Basarir).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

http://dx.doi.org/10.1016/j.ajmsc.2015.12.001

^{1319-5166 © 2016} King Saud University. Production and Hosting by Elsevier B.V. All rights reserved.